



Food and Feeding Habit of Kob (*Kobus kob*) in Kainji Lake National Park

*¹Fingesi, U.I., ²Orsar, J. T. and ²Tyowua, B.T.

¹Department of Wildlife and Ecotourism, FCWM, New Bussa, Niger State, Nigeria

²Department of Wildlife and Range Management, Federal University of Agriculture, PMB 2373, Makurdi, Nigeria

*E-mail of corresponding author: irokau@gmail.com

ABSTRACT

This study focused on the food and feeding habit of Kob (*Kobus kob*) in Kainji Lake National Park. Feed utilization, selection and quality of grasses utilized for food by Kob (*Kobus kob*) were examined. Feed plants and parts grazed by *Kobus kob* were identified using standardized procedures. The procedure initially involved general observations of all group members before focusing on one individual seen to be most stable and easily identified, while grazing between 7:00-12:00 h and 16:00-18:00 h. The results revealed that eighteen plant species were identified as feed components for kob population in the Park in both wet and dry seasons. Wet season forage utilization included *Andropogon gayanus* (23.43%), *Andropogon tectorum* (10.8%) *Gardenia aqualla* and *Grewia mollis* (0.91%) utilization respectively. Dry season forage utilization include *Grewia mollis*(0.91%) and *Azelia africana* (14.40%)utilization. Young leaf material was the most preferred part of plant species (50.88%), followed by Fruits/ Soft seed coat material (19.88%), while Bark of tree was the least preferred part of plant (1.17%) utilized. There was no significant difference ($P>0.05$) in plant species utilized as feed by *Kobus kob* populations in the Park in both wet and dry season. However there was a highly significant difference *** ($P>0.05$) in parts of plants grazed by *Kobus kob* in the park. Illegal activities such as poaching and cattle grazing are found to be prevalent in KLNP, hence measures to improve the feed resources of *Kobus kob* populations in the Park are required. It is therefore recommended that proper monitoring of wildlife resources is crucial, monitoring to ensure compliance by all people so as to minimize grazing and other illegal activities around the park.

Keywords: Food Plants, Season, Kob, Utilized and KLNP.

INTRODUCTION

According to Janean and David (2014) Wildlife foods are generally described by their availability and palatability. Availability refers to the season the food is present in the habitat, how much food there is, and how easily animals can access food. Palatability describes the nutritive and digestible qualities of food. Wildlife selects and eats palatable foods more often than other foods. Different levels of wildlife food preferences are recognized: 1) preferred foods have high nutritional value and are eaten more frequently; 2) moderate value foods, or staples, provide adequate nutrition, but are usually the animal's second choice; and 3) stuffing or emergency foods provide low nutritional value. Wildlife eats these foods in large amounts only when nothing else is available.

Assessing food quality is an important issue when managing and studying herbivores, such as kob. The quality of vegetation relates to the fertility of the soil. The higher the soil fertility, the better the nutritional quality, or palatability, of the vegetation. Many young plant shoots are extremely palatable early in the growing season. The quantity of available food often is the strongest limiting factor in food resources (Waterman and Kool, 1994).

Herbivores rely entirely on plant materials to meet their nutrient and energy requirements (Waterman and Kool, 1994). They face special challenge in selecting optimal diet. Plants and parts are not simply discrete packets of nutrients, they also contain a range of metabolites that are variously refractory to digestion and capable of lowering the efficiency with which nutrients can be obtained or actually harmful to the animal through interference with normal physiological processes (Van Soest 1992).

Kobs are grazers of the commonest grasses, notably *Hyparrhenia*, *Brachiaria*, *Setaria* and *Paspalum* species, among many others. Seasonal change in diet emphasizes the kobs preference for a short sward (Kingdon 1997). Most of daily activity of kob is spent feeding and ruminating with other activities in between; feeding is the dominant activity in early hours and late evening in the day (Agbelusi 1991).

The rate of food intake by herbivores depends on the nutritive quality of the food for example in domestic sheep, intake rate first increases and then decreases as the energy quality of food declines (Fryxell *et al.*, 2014). Grass growth rates in the rainy season are more than sufficient for the animals, but in the dry season available food fell below maintenance requirements.

Food is continuously distributed across the landscape, rather than in definable patches. Animals graze according to marginal value rule, which implies concentrating its feeding in sites where food was higher than average (Distel *et al.* 1995). This lead to the animal having greater opportunity to develop detailed knowledge of the landscape as seen in the cattle (Distel *et al.* 1995).

Nutritional values of plant species

According to Bothma (1996) the nutritional value of plant species is determined by the degree to which the animal eats the plant material voluntarily and the efficiency with which the nutrient can be used for maintenance and plant production. The most valuable plant species are those which are regularly eaten and have a high nutritional value usually at the early growing stages. In the course of the growing season the quantity of food increases while the quality decreases (Bothma 1996). The most important component in food plants for animals is protein (Bothma 1996). It is maintained that only a 5% crude protein level in the vegetation is required for African ungulates and 8% for growing livestock's. This study is aimed at determining the nutritional ecology of *Kobus kob*; identifying food plants species and feeding habits of *kobus kob* in Kainji Lake National Park.

MATERIALS AND METHOD

The Study Area

Kainji Lake National Park is located between latitudes 09°40'N - 10°30'N and longitudes 03° 30'E - 5° 50'E. The park covers a total area of 5340.82sq.km. The Kanji Lake National park was established in 1979 (under decrees 46 of 29th July, 1979), thereby making Kanji lake National park one of the most important National park in Africa, as is highly endowed with many flora and fauna resources. It is made up of two non-contiguous sectors, the Borgu and Zugerma sectors with Borgu sector comprising 3,970.83 km² (74.3%) and the Zugerma sector covering an area of 1,370 km² (25.7%). The vegetation of the Borgu sector has been described as Northern Savanna. The six main vegetation communities in Park are (i) *Burkea africana/Detarium microcarpum woodland savanna* (ii) *Diospyros mespiliformis* dry forest (iii) Riparian forest and woodland (iv) *Terminalia macroptera* tree savanna (v) *Isobertinia tomentosa* woodland and (vi) *Isobertinia doka*, savanna woodland (Ayeni, 2007).

The Oli river flows from the Republic of Benin through Borgu sector into the Niger river. In the dry season, the river breaks into pools that hold water throughout the year and serve as the only source of water for the wild animals. Long term average annual rainfall is between 900 and 1,100 mm. The Park is blessed with diverse fauna resources including *Papio anubis*, *Kobus kob*, *Hippopotamus amphibus*, *Syncerus caffer*, *Panthera leo*, *Panthera pardus*, *Hippotragus equinus* and *Alcelaphus buselaphus* amongst others. Woody resources in the Park include *Burkea africana*, *Terminalia avicennoides*, *Diospyros mespiliformis*, *Anogeissus leiocarpus*, *Entanda africana* and *Vitex doniana* amongst others. Forage species in the Park include *Hyparrhenia dissoluta*, *Andropogon gayanus*, *Brachiaria brachyiopha*, *Hyparrhenia rufa*, *Adropogon perligulatus*, *Andropogon tectorium* and *Beckeropsis uniseta* amongst others (Ayeni *et. al.*, 1982).

Reconnaissance Survey

A Reconnaissance survey was carried out in the study area prior to the detailed study to identify areas where Kobs (*Kobus kob*) are found in Kainji Lake National Park.

Study Design

The design of this study was based on the methodology adopted from Akanbi (1997), which stated that an ecological survey for an area should be conducted on comparative bases, particularly the heterogeneous to indicate a long term range. The study was carried out in the Borgu sector of Kainji Lake National Park. Already existing transects and trails transects was used for this study. Out of a

total of ten (10) existing/ accessible transect in the Borgu sector of Kainji Lake National Park. Five transect was randomly selected for the study in the area. Transects length ranges between 22km to 45km.

Data Collection Techniques

Direct Method of Observation

Kobs were identified as described by Jean and Pierre (1990). Feed plants and parts eaten by Kobs were identified through direct observations with the aid of Zeiss Dialyt Binoculars (10 x 40). Feeding observations followed the scanning focal point techniques as described by Altman (1994). The procedure initially involved general observations of all group members before focusing on one individual seen to be most stable and easily identified. Observations were made on several groups of Kobs that comprised between 4 and 12 individuals from an average distance of 20 m. A camera was used to take pictures of the plants eaten were necessary. Also a field guide book titled “A handbook of west African weeds” by Akobundu and Agyakwa (1989) was used to identify the plant species utilized by kob. The following parameters were recorded as food items species, young leaves, mature leaves and flowers, a total of 192 h of direct observations were made from 20 km transect present in each of the five vegetation communities identified in the Park. Each transect was traversed twice in a month for a period of 24 months between May 2015 and June, 2017.

Feeding observations were usually made between 07:00-12:00 h and 16:00-18:00 h, which was considered to be the peak activity period. Feed items were collected after the group had finished feeding and moved out of the site, a herbarium was prepared from them and later identified as described by Stanfield (1970), Lowe (1989) and Keay (1989).

Data Analysis

Percentage plant and parts utilized as feed was calculated for each plant species as recommended by Alika (2006) as follows:

$$F_k = \sum Y_i / n \times 100$$

- Where, F_k is the frequency for species K, Y_i is the incidence (presence or absence) of species K, n is the total number sampled.

Feed materials were analyzed for proximate, minerals, vitamins and amino acids contents as recommended by AOAC (2000). All data collected were subjected to two ways Analysis of Variance (AVOVA) at (p<0.05) significant level as recommended by Alika (2006).

RESULTS AND DISCUSSION

RESULTS

The result from this study is presented below;

From the study carried out eighteen plant species were identified as feed components for kob population in the Park in both wet and dry seasons. Fourteen plant species were grazed and browsed in the wet season. Wet season forage utilization showed that *Andropogon gayanus* young leaves was the most utilized grass species with (18.02%) utilization, followed by *Gardenia aqualla* (6.31%) and *Grewia mollis* (5.86%) utilization (Table 1), while, the least preferred grass species includes *Acroceras zizanioides* (0.91%). Young leaf material was the most preferred part of plant species (59.45%), followed by matured leave material (26.13%), while bark of tree was the least preferred part of plant (0.91%) utilized.

Dry season forage utilization by kob equally followed the same trend when compared with wet season forage utilization, in which sixteen plant species were utilized as feed by kob, *Hyparrhenia rufa* (12.87%) followed by *Azelia africana* (9.36%) and *Panicum Paucinode* had (7.02%), while the least utilized is the *Acroceras zizanioides* having (0.58%) utilization only. (Table 2). Young leaf material was the most preferred part of plant species (50.88%), followed by Fruits/ Soft seed coat material (19.88%), while bark of tree was the least preferred part of plant (1.17%) utilized.

The result on the Mean seasonal percentage utilization of forage plant species by Kob in Kainji Lake National Park (fig.1) shows that *Andropogon gayanus* is the most preferred food plant in the wet season having (23.43%) preference while *Grewia mollis* (16.38%) is most preferred in the dry season. *Acroceras zizanioides* is the least preferred food plant in both wet and dry seasons having low preference rate of 0.91% and 0.54% respectively.

Proximate composition of major grass species utilized as feed by kob in the Park indicated that *Andropogon gayanus* has a crude protein (7.60%) and fat contents (5.11%), while, crude fibre content is (35.10%) (Table 3).

Table1: Plants and parts utilized as feed in wet season by *Kobus kob* in Kainji Lake National Park

S /no	Types of food eaten	Parts eaten					Part Utilized (%)				
		ML	YL	D L	F/ SC	BT	ML	YL	D L	F/ SC	BT
1	<i>Acroceras zizanioides</i>	0	0	0	x	0	0	0	0.91	0	0
2	<i>Afzelia africana</i>	0	0	0	0	0	0	0	0	0	0
3	<i>Andropogon gayanus</i>	x	x	0	0	0	5.41	18.02	0	0	0
4	<i>Andropogon tectorum</i>	x	x	x	0	0	3.6	4.5	2.7	0	0
5	<i>Cassia mimosoides</i>	0	0	0	0	0	0	0	0	0	0
6	<i>Cyperus difformis</i>	0	x	0	x	0	0	3.6	0	1.8	0
7	<i>Cyperus iria</i>	0	0	0	0	0	0	0	0	0	0
8	<i>Digitaria ciliaris</i>	x	x	0	0	0	3.6	1.8	0	0	0
9	<i>Echinochloa obtusiflora</i>	x	x	0	0	0	3.15	4.5	0	0	0
10	<i>Gardenia aqualla</i>	0	x	0	x	0	0	6.31	0	3.6	0
11	<i>Grewia mollis</i>	x	x	0	0	0	4.05	5.86	0	0	0
12	<i>Hyparrhenia</i>	x	x	0	0	0	2.7	3.15	0	0	0
13	<i>Imperata cylindrical</i>	0	x	0	x	x	0	1.8	0	2.7	0.91
14	<i>Khaya senegalensis</i>	0	0	0	0	0	0	0	0	0	0
15	<i>Maranthes polyandra</i>	x	x	0	0	0	1.8	3.6	0	0	0
16	<i>Panicum Paucinode</i>	x	x	0	0	0	0.91	2.7	0	0	0
17	<i>Paspalum scrobiculatum</i>	x	x	x	0	0	0.91	2.7	1.8	0	0
18	<i>Pterocarpus erinaceous</i>	0	x	0	0	0	0	0.91	0	0	0
							26.13	59.45	5.41	8.1	0.91

Note x: Utilized, 0: Not Utilized ML=Mature leaves, YL =Young leaves, D L = Dry leaves, F/ SC = Fruits/ Soft seed coat, BT = Bark of tree

Table 2: Plants and Parts Utilized As Feed in Dry Season by *Kobus kob* in the Park

Types of food eaten	Parts eaten					Part Utilized (%)				
	ML	YL	D L	F/ SC	BT	ML	YL	D L	F/ SC	BT
<i>Acroceras zizanioides</i>	0		x	0	0	0	0	0.58	0	0
<i>Azelia africana</i>	0	x	0	x	0		9.36		4.68	0
<i>Andropogen gayanus</i>	0	0	0	0	0	0	0	0	0	0
<i>Andropogen tectorum</i>	x	x	x	0	0	4.68	3.51	4.09	0	0
<i>Cassia mimosoides</i>	x	x	x	x	0	2.92	1.75	1.17	2.34	0
<i>Cyperus difformis</i>	0	x	0	x	0	0	0.58	0	0.58	0
<i>Cyperus iria</i>	0	0	0	x	0	0	0	0	1.17	0
<i>Digitaria ciliaris</i>	0	x	0	x	0	0	0.58	0	0.58	0
<i>Echinochloa obtusiflora</i>	0	0	0	0	0	0	0	0	0	0
<i>Grewia mollis</i>	0	x	x	x	0	0	1.17	1.17	2.34	0
<i>Hyparrhenia</i>	x	x	0	0	0	3.51	12.87	0	0	0
<i>Imperata cylindrical</i>	x	x	x	x	0	1.17	3.51	1.17	1.17	0
<i>Khaya senegalensis</i>	0	x	0	x	x	0	2.34	0	3.51	1.17
<i>Maranthes polyandra</i>	0	x	0	x	0	0	4.68	0	2.34	0
<i>Panicum Paucinode</i>	x	x	x	0	0	2.34	7.02	2.92	0	0
<i>Paspalum scrobiculatum</i>	0	x	0	0	0	0	1.17	0	0	0
<i>Pterocarpus erinaceous</i>	0	x	x	x	0	0	1.17	2.34	1.17	0
<i>Seteria. sphacelata</i>	0	x	0	0	0	0	1.17	0	0	0
						14.62	50.88	13.44	19.88	1.17

Note x: Utilized, 0: Not Utilized ML=Mature leaves, YL =Young leaves, D L = Dry leaves, F/ SC = Fruits/ Soft seed coat, BT = Bark of tree




Fig. 1: Mean Seasonal Percentage Utilization of Forage Plant Species by Kob in Kainji Lake National Park

Table 3: Proximate Composition of Major Feed Items Eaten by *Kobus kob* Population in the Park

Plant species	DM	CP	CF	FAT	ASH	NFE
1. <i>Andropogon gayanus</i>	33.5	7.60	35.10	5.11	9.42	42.77
2. <i>Andropogon tectorum</i>	39.4	6.26	32.11	4.77	8.65	48.21
3. <i>Cassia mimosoides</i>	31.3	6.10	30.00	3.09	5.67	30.43
4. <i>Echinochloa obtusiflora</i>	48.3	7.21	42.31	4.91	7.42	38.15
5. <i>Hyparrhenia rufa</i>	39.1	9.03	32.50	4.11	8.12	46.17

Adopted from Aremu and Onadeko (2008) : %DM: Percentage dry matter, %CP: Percentage crude protein, %CF: Percentage crude fibre, %NFE: Percentage nitrogen free extract

DISCUSSION

The findings of the study revealed that the frequencies of feeding on *Andropogon gayanus* in wet season and *Andropogon tectorum* in wet and dry seasons by kobs explains why kobs are mostly grazers. *Grewia mollis*, *Maranthus polyandra*, and *Azelia africana* were fed upon by kobs mostly during the dry season as browse plants. Agbelusi (1989) asserted that kob utilizes both browse and grass species and that the grass species favoured by kob were mainly perennials thus making it possible for them to select herbaceous plants that are available all the year round.

There was no significant difference ($P > 0.05$) in plant species utilized as feed by *kobus kob* populations in the Park in both wet and dry season. Nevertheless there was a highly significant difference *** ($P > 0.05$) in parts of plants grazed by *kobus kob* in the park.

Feed Selection and the Nutritional Values of Plant Species

The most preferred part of plants in both wet and dry seasons is the young leaf (59.45%) and (50.88%), respectively when compared to matured leaf (26.13 and 14.62%), respectively. This may

not be unconnected to the fact that young leaves are more palatable, succulent and easy to digest making the feed nutrient easily available to animal for growth, maintenance of pregnancy and lactation (Voeten, 1999). Kobs only nib at the tip of the plant species; this may explain why ungulate preferred new flush of grasses from newly burnt rangelands where it can easily cut the leaf tip (Mwangi *et al.*, 1999; Aremu, 2001). Mature leaves were less preferred by kobs population due to the fact that they are dry and coarse thereby hindering digestibility. These observations followed that of Holechek (1994) and Stoddart *et al.* (1995). Kobs in the park show grazing preference for swards of intermediate grass height and biomass, they are mixed feeders, changing from grazing in the rains to browsing in the dry season. Their habitats are varied and ranges from dense woodland and to open flood plains. These observations also followed that of (Fryxell *et al.* 2004). *Andropogon gayanus* was the most preferred grass species in the rainy season due to the fact that it contains high percentages of crude protein and fat 7.60 and 5.11%, respectively, which is above the 5% crude protein level in the vegetation that is required for African ungulates growth (Bothma 1996). While other grasses have high level of CP and lower fat content.

CONCLUSION

Frequent bush burning which has quite an adverse effect on the plant especially grass growth and other illegal activities such as poaching and cattle grazing are found to be prevalent in KLNP. For the park to increase the habitat potentials and to improve grazing opportunities for the *Kobus kob* population in the Park, most especially during dry season when the habitat resources are usually over stretched. There is therefore, the need to improve habitats quality through application of controlled burning programmes, planting of desirable grasses and prohibit illegal grazing by livestock within Park boundaries, to eliminate competition for limited habitat resources between wild and domestic animals.

ACKNOWLEDGEMENTS

Author is thankful to research supervisors; Orsar, J. T. and Tyowua, B.T. for their valuable advice, encouragement and guidance throughout the study period. The author is highly indebted to the Conservator General National Park Service Abuja, Nigeria for granting permission to conduct research at the National Park.

Author's gratitude goes to the Conservator of Park, KLNP, Mr. Kabiru Mohammed for his support and encouragement during the period of the research at the National Park. Finally author appreciate the tireless efforts of the technical assistant of the FCWM/KLNP Park Staff; Ishaya T. Dung, Yakubu Abu, Sunday Akor, Hassan Bashiru, and Mutiu.

REFERENCES

- Agbelusi, A. (1989). Feeding-habits of the Senegal Kob (*Kobus kob kob*, Erxeleben 1777) under Ranching conditions and in the wild. *Applied Animal Behaviour*, 23: 179-185.
- Agbelusi, A. (1991). Variations in activities of Senegal Kob (*Kobus kob kob*, Erxeleben 1777) during daylight period in an enclosure at Kainji Lake Research Institute range farm, New Busa, Nigeria. *Afr. J. Ecol.* 29: 353-355.
- Akanbi, O. A. (1997): An Ecological Basis For Management of Enamania Game Reserve in Benue State, Nigeria. Ph D. Thesis unpublished, Department of Wildlife and Fisheries, University of Ibadan.
- Akobundu, I.O. and Agyakwa, C.W. (1989). A handbook of West African weeds, IITA, Ibadan, pp546.
- Alika, J.E., (2006). Statistics and Research Methods. 2nd Edn., AMBIK Press, Benin City, Pages: 366.
- Altman, J., (1994). Observational study of behaviour sampling methods. *Behaviour*, 49: 257-267.
- AOAC, (2000). Association of Official Analytical Chemists. Official Methods of Analysis, Washington DC.
- Aremu, O.T., (2001). Effect of fire on plants, soil nutrients and soil micro-organization of Old Oyo National Park, Nigeria. *J. Agric. For. Fish.*, 2: 36-40.

- Aremu, O.T. and Onadoko, S.A. (2008). Nutritional Ecology of African Buffalo (*Syncerus caffer nanus*). *International Journal of Agricultural Research*, 3: 281-286
- Ayeni J.O.S, Afolayan TA and Ajayi SS (1982). Introductory handbook on Nigeria wildlife: Kainji Lake Research Institute. New Bussa, Nigeria, pp 43 – 45.
- Ayeni, J.S.O. (2007). Participatory Management Plan of Kainji Lake National Park: Produced by Natural Resources, Agricultural and Environment Consultants. Environ- Consult. Lagos Pg. 150.
- Bothma, J .du P. (1996). Game ranch management. 3rd edition. J.L. van Schaik Publishers, 1064 Acradia Street, Hatfield, Pretoria. Pp.165-402
- Distel, R.A., Laca , E. A., Griggs., T.C., and Dement, M.W.(1995). Patch Selection by Cattle : Maximization Of Intake Rate In Horizontally Heterogeneous Pastures. *Applied Behavior Science* 45: 11-21.
- Fryxell, J. M., Wilmshurst, J. F., Sinclair, A. R. E., and (2004). Predictive models of movement by Serengeti grazers. *Ecology* 85: 2429- 2435.
- Fryxell, J.M.,Sinclair , A.R.E.,and Caughley G (2014). Wildlife ecology, conservation and management. Third ed. WILEY Blackwell. Pub. By JOHN Willey and sons ltd.
- Laserwords Private Limited , Singapore Markono Print Media Pte Ltd. Chennai, India.pp 26-279.
- Holechek, J.L., (1994). Comparative contribution of grasses, forbs and shrubs to the nutrition of ungulates. *Rangelands*, 16: 645-648.
- Janean H. Creighton, and ., David M. (2014) Baumgartner Wildlife Ecology and Forest Habitat. EB1866. WSU Cooperative Extension. Pullman, *Washington*
- Jean, D. and D. Pierre, 1990. A Field Guide to the Large Mammals of Africa. 3rd Edn., Collins St. James Publication, London, pp: 401
- Keay (1989). Tress of Nigeria. 1st Edn., Oxford Science Publication, New York, pp: 476.
- Kingdon, J. (1997) The Kingdom field guide to African mammals. Natural world academic press Harcourt Brace and company publisher San Diego London pp403-404.
- Lowe, J., (1989). The Flora of Nigeria Grasses. 2nd Edn., Ibadan University Press, Nigeria pp: 326.
- Mwangi, D.M., G. Calisch, W. Thorpe and K.E. Giller, (1999). Harvesting management option for legumes intercropped in Napier grass in the central highlands of Kenya. *Trop. Grasslands*, 37: 94-102.
- Stanfield, D.P., (1970). The Flora of Nigeria Grasses. 1st Edn., Ibadan University Press, Nigeria.
- Stoddart, L.A., A.D. Smith and T.W. Box, 1995. Range Management. Mc Graw Hill Book Co., New York, pp: 532.
- Van Soest, P.J., (1992). Nutritional Ecology of Ruminants. 2nd Edn., O and S Books, Corvallis OR.
- Voeten, M.M., (1999). Co-existence of wildlife and livestock in East African savanna system. *Trop. Resour. Manage.*, 29: 84-109.
- Waterman, P.G. and K.M. Kool, 1994. Food Selection and Plant Chemistry. In: Animal Food Requirement, Davies, A.G. and J.E.O. Oats (Eds.). University Press, Cambridge, London.